Published online 2017 June 30.

Research Article

Musculoskeletal Related Injuries After 2017 Kermanshah Earthquake

SM Javad Mortazavi¹, Mohammad Ebrahiminasab¹, Amir Reza Farhoud ¹, Babak Mirzashahi¹, Sadegh Saberi¹ and Ehsan Ghadimi^{1,*}

¹Orthopedics Department, Joint Reconstruction Research Center, Tehran University of Medical Sciences, Tehran, IR Iran

[°] *Corresponding author*: Ehsan Ghadimi, Keshavarz Blvd, Gharib Street, Imam Khomeini Hospital Complex, Joint Reconstruction Research Center (JRRC), Postal code: 1419733141, Tehran, IR Iran. Tel: +98-9128415815, E-mail: ehsan.ghadimie@gmail.com

Received 2017 April 23; Revised 2017 May 15; Accepted 2017 June 22.

Abstract

Objectives: This study aims to investigate the pattern of Musculoskeletal and soft tissue injuries in a series of referred patients in a tertiary medical center and review previous reports in the literature.

Methods: In this study, we recorded musculoskeletal injuries of the victims, following the Kermanshah earthquake on November 12, 2017, with a total number of 38. All patients were admitted to the Imam Khomeini Hospital. Injury type, site of injury, gender, age, complications, mortality, and treatment modality were recorded for each case.

Results: Among 38 patients, 18 were male (47%) and 20 were female (53%). The mean age was 37 ± 21.6 years. Most of them were aged between 18 and 65 (71.1%). Lower extremities were involved more than the upper (37% versus 24%) and distal limb involvement was more than the proximal (57.3% versus 43.7%). One patient died due to necrotizing fasciitis and septic shock after femoral shaft open fracture. Two patients developed compartment syndrome and underwent fasciotomy complications by skin necrosis and infection. **Conclusions:** Soft tissue injuries and fractures are the most common injuries following the earthquakes, hence orthopedic surgeries play a vital role in disasters. Despite patients overload and emergency situation in natural disasters like earthquakes, it is crucial to have a stepwise and evidence based approach for each patient. For the patients with open fractures referred to a tertiary center, careful contamination assessment even in the previously managed wounds is highly recommended, especially for those with early wound closure. Early fasciotomy in the crush syndrome is not beneficial as it has adverse effects such as skin necrosis and infection.

Keywords: Earthquake, Disasters, Iran, Orthopedic Procedures, Musculoskeletal Injuries

1. Background

Natural disasters lead to enormous loss of life and economical forfeiture (1). Tsunamis and floods result in higher mortality, whereas higher morbidity occurs as a result of earthquakes, mostly orthopedics (1-3). Considered as one of the most frequent natural calamities, there were at least 8000 deaths and 26,000 injuries due to the earthquakes each year (4). After the management of the life-threatening injuries, the leading causes of morbidity are assumed to be soft tissue and extremities injuries (5-7). On November 12, 2017, at 21:18 p.m. local time, Kermanshah, Iran, the region of Sarpol-e Zahab, was struck by a 7.3-Richter earthquake. As the earthquakes affect a great deal of victims, better epidemiological knowledge of the injuries is essential for improving the disaster management (8-13). This study aims to investigate the pattern of the Musculoskeletal and soft tissue injuries in a series of referred patients in a tertiary medical center.

2. Methods

In this study, we recorded musculoskeletal injuries of the victims, a total number of 38, who were transported by airplane to our hospital, a tertiary medical center (Imam Khomeini Hospital). Injury type, site of injury, gender, age, complications, mortality, and treatment modality were recorded for each case. All of the patients received first aid treatments. The individuals who were candidates for surgery received pre-operative antibiotics. Strict glycemic control was performed for the patients with a history of Diabetes Mellitus. The final diagnosis of the injuries was made by the hospital physicians, based on clinical signs, symptoms, and necessary imaging.

3. Results

A total of 630 people died and more than 7,000 were injured as a result of the earthquake (14). More than half of the Iranian casualties were from Sarpol-e-Zahab and the

Copyright © 2017, Journal of Orthopedic and Spine Trauma. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

Ezgeleh district, which had a population of over 30,000. At least 7 people died and another 500 injured in the adjacent area in Iraq, according to their officials (15). About 70,000 individuals, from 14 Iranian provinces, became homeless. The earthquake completely destroyed around 12,000 houses and damaged another 15,000 (16).

In total, 38 injured patients were admitted to our hospital on November 13, 2017. Of the patients, 18 patients were male (47%) and 20 patients were female (53%). The mean age was 37 ± 21.6 years. A total of 18.4% of the patients were under the age of 18, 71.1% between 18 and 65, and 10.5% above 65 years. There were more females than males in patients below 65 years of age (Figure 1). Spine, lower, and upper limb involvement was noted in 6 (16%), 14 (37%), and 9 (24%) of the individuals. Of the patients, 7 (18%) of them had pelvic fracture and 8 (21%) had no fractures (Table 1). Proximal limb involvement (Femur and Humerus) was noted in 7 (43.7%) and distal limb involvement (Tibia, Ulna and Radius) was noted in 16 (57.3%) of the patients. Acute renal failure emerged in 1 patient (2.6%).



Figure 1. Age Groups and Gender Distribution

One patient (2.6%) died due to septic shock, he was a 30year-old man who had femoral shaft open fracture. The initial measures including wound approximation were taken before admission in this center. The vital signs became unstable in the 1st hours; the blood pressure was about 66/55 mmHg and the heart rate 150 Bpm. The clinical examinations were assumed to be in favor of hypovolemic shock, thus, Aggressive Resuscitation was performed, however, there were no ameliorations and the patient intubated within the next few hours. A total of 15 hours later, the paTable 1. Distribution of the Fractures and the Treatment Based on the Anatomical Sites

Localization	No. (%)	Number of Surgeries	Number of Conservative Therapies
Lower limb fractures			
Femur	5 (35)	5	
Foot	4 (29)	3	1
Ankle	4 (29)	4	
Leg	1(7)	1	
Upper limb fractures			
Scapula	1 (12.5)		1
Arm	1(12.5)	1	
Forearm	1 (12.5)	1	
Wrist	2 (25)	2	
Elbow	1(12.5)	1	
Hand	3 (37.5)	3	
Spine	6 (16)	2	4
Pelvic	7 (18)	2	5

tient underwent surgery for temporary stabilization with external fixator. Massive Purulent discharge was noted and as the wound explored, extensive necrotizing fasciitis and myonecrosis were seen. The wound was probably contaminated by the patients own fecal. Being in septic shock and fasciitis extending to the proximal of the lower limb, hip disarticulation was performed. Unfortunately, Later aggressive irrigation and debridement's were not successful, and the patient died 48 hours later.

There were 2 patients (5.26%) that experienced compartment syndrome. Early fasciotomy was done for both of them. Both were complicated by an infection and stayed in the hospital for more than 1 month.

Two patients (5.26%) experienced cervical spine fracture dislocations. Both of them underwent early surgery within 12 hours. One of the patients who had normal neurologic examination prior to surgery, fully recovered after surgery and discharged 3 days later. However, the other one who had complete paraplegia prior to surgery did not recover.

One patient experienced paraplegia without a fracture or cord injury in neither the CT scan nor the MRI of the spine. He fully recovered gradually and was discharged 1 week later. The diagnosis was assumed to be the crush syndrome of the spine with no structural abnormality.

4. Discussion

Earthquakes are one of the most fatal mass impact humanitarian disasters, causing a high mortality and morbidity each year. Population density of the region, time of day of the earthquake, and building infrastructure quality are the factors determining the extent of the damage (17). Similar to us, Kaim Khani et al. noted that young people, especially females, are more susceptible for orthopedic injuries (18). On average, pediatrics are affected more, both in developing and developed countries (19, 20); however, Sarisozen and Durak stated that compared to the adults, pediatrics have lower amounts of injuries but the same severity (21).

4.1. Percentage of Orthopedic Injuries

Limb injuries are considered as the most common injury, accounting for 54% - 66% (22-24). A total of 77% of the total victims from the Bam earthquake in Iran (25), 84% in Van earthquake in Turkey (26), 88% in Wenchuan earthquake in China (27), 89% in Battagram in Pakistan (18), and Gujarat earthquake in India (28) had orthopedic injuries. In our study, 8 (21%) of the patients had no fractures and 10f the injured patients had a head injury. The other 29 patients (76%) had orthopedic related injuries, similar to previous reports.

Fractures, crush injuries, and soft tissue injuries are considered to be the most common type of orthopedic-related injuries (25-27, 29).

4.2. Anatomic Distribution

Following the 2001 India earthquake, among 281 orthopedic injuries, lower limbs were affected the most with 204 (72%), followed by upper limbs 57 (20%), and pelvis 20 (8%) (30) injuries. Similar results from Iran (25, 31), Turkey (26, 29), Haiti (32, 33), Nepal (34, 35), China (27, 36), and Pakistan (18, 37) were reported, demonstrating high prevalence of the lower limb fractures, followed by upper limbs, and pelvis. The proximal bone involvement is greater if the earthquake happens early in the morning (as most people are asleep) and it is assumed that proximal bones are more involved generally (25).

In our study similar results were found, lower limbs affected 37%, upper limbs 24%, and pelvis 18%. According to the previous studies, pelvic injuries include a small portion of the injuries, which may be due to the fact that patients died before reaching the hospital (38). In our study 7 people (18%) had a pelvic injury, 2 of them underwent surgery, and conservative therapy was chosen as the treatment for the rest of them.

4.3. Orthopedic Procedures

Orthopedic procedures following an earthquake include debridement, external fixation, open reduction, and internal fixation (ORIF), close reduction and casting, as well as amputation. Debridement is the most common type of procedure (26, 29, 32), however, there is a significant difference in the other treatments in previous reports. Guner et al. reported (29) the greatest number of patients who received orthopedic procedures. Among 370, the most common procedure was debridement (45%), followed by ORIF (24%), CR and casting (20%), external fixation (8%), and amputation (3%). Phalkey et al. (30) reported debridement in 26%, close reduction and casting in 30%, ORIF in 28%, external fixation in 1%, and amputation in 11% of the patients. Bar-On et al. (32) reported debridement in 68% of cases followed by external fixation in 55% of cases. In our study, 4 patients (13%) underwent debridement, 20 patients (66%) ORIF, 1 case (3%) needed external fixation, and 1 (3%) patient underwent amputation.

The importance of the limb salvage surgeries, in the situations other than disasters, is obvious (39). It matters more when a natural calamity happens in a poorer economy (40). Despite good rehabilitations after the amputation (41), the individuals will encounter financial problems following that (40, 42).

4.4. Open Versus Closed Fracture

Open fractures usually occur after the earthquakes, however, not more than closed fractures. Open fractures occur in 8% - 13% of the affected people (27, 30). In the Pakistan earthquake in 2011 open fractures were reported as 35% and closed fractures 65% (18). In Nepal (34), Haiti (32), China (27), and Turkey (26, 29) open fractures were 27%, 28%, 34%, and 14%.

Similarly, in our experience, only 4 (12%) open fractures were seen. One of them complicated by necrotizing fasciitis, which led to death and 2 of them complicated by skin necrosis and infection. One patient underwent external fixation without any complications.

4.5. Crush Syndrome and Compartment Syndrome

Among 228 orthopedic-related injuries, Tahmasebi et al. (25) reported 6 patient with (2%) crush syndrome and 18 (6%) patients with compartment syndrome. In China earthquake, Dai et al. (27) reported 19 cases (7%) of crush syndrome and 18 cases (6.5%) of compartment syndrome.

Crush syndrome among pediatric patients was reported by Iskit et al. and Donmez et al. (43, 44). Sarisozen reported acute renal failure as a result of Crush syndrome (21). In our patients, compartment syndrome happened in 2 of them and 1 of the patients had acute renal failure, none

of them were among the pediatric group. Both of the patients who underwent fasciotomy were complicated by infection. Due to the fact that in crush injuries the muscles are already dead, fasciotomy is not indicated to save at-risk muscles like other forms of compartment syndrome. As previously noted, routine fasciotomies does not only not improve outcomes, but it also increases the risk of renal failure, wound infection, and sepsis related to wound infection (45-48).

4.6. Infection

The range of infection is estimated to be 19% - 35% (49, 50). Risk factors for infection include multiple bone fracture, crush syndrome, amputation, and paraplegia (51-53). Infection was seen in 3 patients in our experience. All of them as a complication of open fracture; 2 patients were treated successfully with antibiotics, irrigation, as well as debridement and skin graft. The last patient with femoral shaft open fracture complicated by necrotizing fasciitis died due to septic shock.

As previously discussed (54, 55), early removal of foreign bodies and organic contaminations (56) accompanied by early irrigation and debridement in the first 24 hours play a crucial role in improving the outcome of open fractures (57, 58). Our unpleasant experience throws the light on this fact that if the wound has been primary closed or approximated, wound reassess mentation is important. Careful patient examination should be done due to contamination possibility. In addition, primary closure of the wounds in open fractures should be done after precise physical examination and careful assessment. Our patients' wound was contaminated by fecal and primary closure was contraindicated (59). Opening the wound, removal of organic contamination, and early and aggressive irrigation and debridement could probably save our patients' life.

Earthquake- related injuries cause long-term morbidities and quality of life decrement (60). Preventing mortality and decreasing morbidities, adequate and short-time aid is crucial (50, 61), in this regard, preparation and planning is suggested for a better disaster management and avoiding further mortality and morbidity. Moreover, it is crucial to have a stepwise and evidence based approach for each patient. For the patients with open fracture referred to a tertiary center, careful contamination assessment, even in previous managed wounds, is highly recommended, especially for those with early wound closure. Besides, not having proven advantages, early fasciotomy in crush syndrome is not suggested.

References

- Seaman J. Disaster epidemiology: or why most international disaster relief is ineffective. *Injury*. 1990;21(1):5–8. discussion 15-6. [PubMed: 2347634].
- Noji EK. Public health issues in disasters. Crit Care Med. 2005;33(1 Suppl):S29–33. [PubMed: 15640676].
- Jiang J, Xu H, Liu H, Yuan H, Wang C, Ye J. Anaesthetic management under field conditions after the 12 May 2008 earthquake in Wenchuan, China. *Injury*. 2010;41(6):e1-3. doi: 10.1016/j.injury.2009.07.071. [PubMed: 19695567].
- Alexander D. Death and injury in earthquakes. *Disasters*. 1985;9(1):57– 60. doi: 10.1111/j.1467-7717.1985.tb00911.x. [PubMed: 20958593].
- Bartels SA, VanRooyen MJ. Medical complications associated with earthquakes. *Lancet.* 2012;**379**(9817):748–57. doi: 10.1016/S0140-6736(11)60887-8. [PubMed: 22056246].
- Redmond AD, Li J. The UK medical response to the Sichuan earthquake. *Emerg Med J.* 2011;28(6):516–20. doi: 10.1136/emj.2009.089920. [PubMed: 20817662].
- Wolf Y, Bar-Dayan Y, Mankuta D, Finestone A, Onn E, Morgenstern D, et al. An earthquake disaster in Turkey: assessment of the need for plastic surgery services in a crisis intervention field hospital. *Plast Reconstr Surg.* 2001;**107**(1):163–8. discussion 169-70. [PubMed: 11176618].
- Eberhart-Phillips JE, Saunders TM, Robinson AL, Hatch DL, Parrish RG. Profile of mortality from the 1989 Loma Prieta earthquake using coroner and medical examiner reports. *Disasters*. 1994;18(2):160–70. [PubMed: 8076160].
- Kaneda M. Injury distributions produced by natural disasters. Asian Med J. 1994;37(10):557–63.
- Noji EK, Kelen GD, Armenian HK, Oganessian A, Jones NP, Sivertson KT. The 1988 earthquake in Soviet Armenia: a case study. *Ann Emerg Med*. 1990;**19**(8):891–7. [PubMed: 2142590].
- Kuthy-Porter J. Earthquake: its consequences in the care of chest diseases in Mexico. *Chest.* 1986;**90**(1):119. [PubMed: 3720372].
- Roces MC, White ME, Dayrit MM, Durkin ME. Risk factors for injuries due to the 1990 earthquake in Luzon, Philippines. *Bull World Health Organ*. 1992;**70**(4):509–14. [PubMed: 1394785].
- Sheng ZY. Medical support in the Tangshan earthquake: a review of the management of mass casualties and certain major injuries. J Trauma. 1987;27(10):1130-5. [PubMed: 3312621].
- IRNA. Western Iran fatal quake death toll rises to 620. 2017. Available from: http://www.irna.ir/en/News/82759872.
- Reliefweb. Iraqi Red Crescent: Increasing the number of the dead to 10 people because of the earthquake and set up camps to shelter the affected families in Darbandikhan and Halapcha. 2017. Available from: https://reliefweb.int/report/iraq/iraqi-red-crescent-increasingnumber-dead-10-people-because-earthquake-and-set-camps.
- Time. Earthquake. 2017. Available from: http://time.com/5020880/ iran-iraq-earthquake/.
- Ramirez M, Peek-Asa C. Epidemiology of traumatic injuries from earthquakes. *Epidemiol Rev.* 2005;27:47–55. doi: 10.1093/epirev/mxi005. [PubMed: 15958426].
- Kaim Khani GM, Baig A, Humail M, Memon M, Quarashi MA. Musculoskeletal injuries among victims of the Battagram, Pakistan earthquake in October 2005. *Prehosp Disaster Med*. 2012;27(5):489–91. doi: 10.1017/S1049023X12001161. [PubMed: 22985734].
- Gnauck KA, Nufer KE, LaValley JM, Crandall CS, Craig FW, Wilson-Ramirez GB. Do pediatric and adult disaster victims differ? A descriptive analysis of clinical encounters from four natural disaster DMAT deployments. *Prehosp Disaster Med*. 2007;22(1):67-73. [PubMed: 17484366].
- Helminen M, Saarela E, Salmela J. Characterisation of patients treated at the Red Cross field hospital in Kashmir during the first three weeks of operation. *Emerg Med J.* 2006;23(8):654–6. doi: 10.1136/emj.2006.036012. [PubMed: 16858109].

- Sarisozen B, Durak K. Extremity injuries in children resulting from the 1999 Marmara earthquake: an epidemiologic study. J Pediatr Orthop B. 2003;12(4):288–91. doi: 10.1097/01.bpb.0000049581.53117.65. [PubMed: 12821848].
- Clover AJ, Jemec B, Redmond AD. The extent of soft tissue and musculoskeletal injuries after earthquakes; describing a role for reconstructive surgeons in an emergency response. *World J Surg.* 2014;**38**(10):2543-50. doi: 10.1007/s00268-014-2607-y. [PubMed: 24791946].
- Peek-Asa C, Kraus JF, Bourque LB, Vimalachandra D, Yu J, Abrams J. Fatal and hospitalized injuries resulting from the 1994 Northridge earthquake. *Int J Epidemiol*. 1998;27(3):459–65. [PubMed: 9698136].
- Mohebbi HA, Mehrvarz S, Saghafinia M, Rezaei Y, Kashani SM, Naeeni SM, et al. Earthquake related injuries: assessment of 854 victims of the 2003 Bam disaster transported to tertiary referral hospitals. *Prehosp Disaster Med*. 2008;23(6):510–5. [PubMed: 19557966].
- Tahmasebi MN, Kiani K, Mazlouman SJ, Taheri A, Kamrani RS, Panjavi B, et al. Musculoskeletal injuries associated with earthquake. A report of injuries of Iran's December 26, 2003 Bam earthquake casualties managed in tertiary referral centers. *Injury*. 2005;**36**(1):27–32. doi: 10.1016/j.injury.2004.06.021. [PubMed: 15589909].
- Gormeli G, Gormeli CA, Güner S, Ceylan MF, Dursun R. The clinical profile of musculoskeletal injuries associated with the 2011 Van earthquake in Turkey. Spine. 2012;28:9.
- 27. Dai ZY, Li Y, Lu MP, Chen L, Jiang DM. Clinical profile of musculoskeletal injuries associated with the 2008 Wenchuan earthquake in China. *Ulus Travma Acil Cerrahi Derg.* 2010;**16**(6):503–7. [PubMed: 21153941].
- Roy N, Shah H, Patel V, Bagalkote H. Surgical and psychosocial outcomes in the rural injured-a follow-up study of the 2001 earthquake victims. *Injury*. 2005;36(8):927–34. doi: 10.1016/j.injury.2005.02.017. [PubMed: 15979621].
- Guner S, Guner SI, Isik Y, Gormeli G, Kalender AM, Turktas U, et al. Review of Van earthquakes form an orthopaedic perspective: a multicentre retrospective study. *Int Orthop.* 2013;**37**(1):119–24. doi: 10.1007/s00264-012-1736-x. [PubMed: 23232655].
- Phalkey R, Reinhardt JD, Marx M. Injury epidemiology after the 2001 Gujarat earthquake in India: a retrospective analysis of injuries treated at a rural hospital in the Kutch district immediately after the disaster. *Glob Health Action*. 2011;4:7196. doi: 10.3402/gha.v4i0.7196. [PubMed: 21799668].
- Emami MJ, Tavakoli AR, Alemzadeh H, Abdinejad F, Shahcheraghi G, Erfani MA, et al. Strategies in evaluation and management of Bam earthquake victims. *Prehosp Disaster Med.* 2005;20(5):327-30. [PubMed: 16295170].
- Bar-On E, Lebel E, Kreiss Y, Merin O, Benedict S, Gill A, et al. Orthopaedic management in a mega mass casualty situation. The Israel Defence Forces Field Hospital in Haiti following the January 2010 earthquake. *Injury*. 2011;42(10):1053–9.
- 33. Blumberg N, Lebel E, Merin O, Levy G, Bar-On E. Skeletal injuries sustained during the Haiti earthquake of 2010: a radiographic analysis of the casualties admitted to the Israel Defense Forces field hospital. *Eur J Trauma Emerg Surg.* 2013;**39**(2):117–22.
- Bar-On E, Blumberg N, Joshi A, Gam A, Peyser A, Lee E, et al. Orthopedic Activity in Field Hospitals Following Earthquakes in Nepal and Haiti. World J Surg. 2016;40(9):2117–22.
- Vaishya R, Agarwal AK, Vijay V, Hussaini M, Singh H. Surgical management of musculoskeletal injuries after 2015 Nepal earthquake: our experience. *Cureus*. 2015;7(8).
- Xiang B, Cheng W, Liu J, Huang L, Li Y, Liu L. Triage of pediatric injuries after the 2008 Wen-Chuan earthquake in China. *J Pediatr Surg.* 2009;44(12):2273–7.
- 37. Bozkurt M, Ocguder A, Turktas U, Erdem M. The evaluation of trauma patients in Turkish Red Crescent Field Hospital following the Pakistan earthquake in 2005. *Injury*. 2007;**38**(3):290–7.
- Calder J, Mannion S. Orthopaedics in Sri Lanka post-tsunami. J Bone Joint Surg Br. 2005;87(6):759–61. doi: 10.1302/0301-620X.87B6.16482.

[PubMed: 15911654].

- Chung KC, Saddawi-Konefka D, Haase SC, Kaul G. A cost-utility analysis of amputation versus salvage for Gustilo type IIIB and IIIC open tibial fractures. *Plast Reconstr Surg.* 2009;**124**(6):1965–73. doi: 10.1097/PRS.0b013e3181bcft56. [PubMed: 19952652].
- Andersson N, da Sousa CP, Paredes S. Social cost of land mines in four countries: Afghanistan, Bosnia, Cambodia, and Mozambique. *BMJ*. 1995;**311**(7007):718–21. [PubMed: 7549685].
- Hettiaratchy SP, Stiles PJ. Rehabilitation of lower limb traumatic amputees: the Sandy Gall Afghanistan Appeal's experience. *Injury*. 1996;**27**(7):499–501. [PubMed: 8977837].
- Walsh NE, Walsh WS. Rehabilitation of landmine victims-the ultimate challenge. *Bull World Health Organ*. 2003;81(9):665–70. [PubMed: 14710508].
- Iskit SH, Alpay H, Tugtepe H, Ozdemir C, Ayyildiz SH, Ozel K, et al. Analysis of 33 pediatric trauma victims in the 1999 Marmara, Turkey earthquake. J Pediatr Surg. 2001;36(2):368–72. doi: 10.1053/jpsu.2001.20719. [PubMed: 11172437].
- Donmez O, Meral A, Yavuz M, Durmaz O. Crush syndrome of children in the Marmara Earthquake, Turkey. *Pediatr Int.* 2001;43(6):678–82. [PubMed: 11737748].
- 45. Sever MS, Erek E, Vanholder R, Akoglu E, Yavuz M, Ergin H, et al. Clinical findings in the renal victims of a catastrophic disaster: the Marmara earthquake. *Nephrol Dialysis Transplant*. 2002;**17**(11):1942–9.
- Nadjafi I, Atef MR, Broumand B, Rastegar A. Suggested guidelines for treatment of acute renal failure in earthquake victims. *Renal Failure*. 1997;19(5):655-64.
- Reis ND, Michaelson M. Crush injury to the lower limbs. Treatment of the local injury. J Bone Joint Surg. 1986;68(3):414–8.
- Reis ND, Better OS. Mechanical muscle-crush injury and acute muscle-crush compartment syndrome: with special reference to earthquake casualties. *Bone Joint J.* 2005;87(4):450–3.
- Keven K, Ates K, Sever MS, Yenicesu M, Canbakan B, Arinsoy T, et al. Infectious complications after mass disasters: the Marmara earthquake experience. Scand J Infect Dis. 2003;35(2):110–3. [PubMed: 12693561].
- Aylwin CJ, Konig TC, Brennan NW, Shirley PJ, Davies G, Walsh MS, et al. Reduction in critical mortality in urban mass casualty incidents: analysis of triage, surge, and resource use after the London bombings on July 7, 2005. *Lancet*. 2006;**368**(9554):2219–25. doi: 10.1016/S0140-6736(06)69896-6. [PubMed: 17189033].
- Huang KC, Lee TS, Lin YM, Shu KH. Clinical features and outcome of crush syndrome caused by the Chi-Chi earthquake. J Formos Med Assoc. 2002;101(4):249–56. [PubMed: 12101860].
- Rathore MF, Rashid P, Butt AW, Malik AA, Gill ZA, Haig AJ. Epidemiology of spinal cord injuries in the 2005 Pakistan earthquake. *Spinal Cord*. 2007;45(10):658–63. doi: 10.1038/sj.sc.3102023. [PubMed: 17228354].
- 53. Tauqir SF, Mirza S, Gul S, Ghaffar H, Zafar A. Complications in patients with spinal cord injuries sustained in an earthquake in Northern Pakistan. *J Spinal Cord Med.* 2007;**30**(4):373–7. [PubMed: 17853661].
- Guthrie HC, Clasper JC. Historical origins and current concepts of wound debridement. J R Army Med Corps. 2011;157(2):130–2. [PubMed: 21805760].
- Kocher MS. Early limb salvage: open tibia fractures of Ambroise Pare (1510-1590) and Percivall Pott (1714-1789). World J Surg. 1997;21(1):116–22. [PubMed: 8943188].
- Robson MC, Duke WF, Krizek TJ. Rapid bacterial screening in the treatment of civilian wounds. *J Surg Res.* 1973;14(5):426–30. [PubMed: 4574478].
- 57. Crowley DJ, Kanakaris NK, Giannoudis PV. Debridement and wound closure of open fractures: the impact of the time factor on infection rates. *Injury*. 2007;**38**(8):879–89.
- Crowley DJ, Kanakaris NK, Giannoudis PV. Irrigation of the wounds in open fractures. J Bone Joint Surg Br. 2007;89(5):580–5. doi: 10.1302/0301-620X.89B5.19286. [PubMed: 17540739].

- Rajasekaran S, Dheenadhayalan J, Babu JN, Sundararajan SR, Venkatramani H, Sabapathy SR. Immediate primary skin closure in type-III A and B open fractures: results after a minimum of five years. *J Bone Joint Surg Br.* 2009;**91**(2):217–24. doi: 10.1302/0301-620X.91B2.21228. [PubMed: 19190057].
- 60. Sudaryo MK, Endarti AT, Rivany R, Phalkey R, Marx M; Besral, et al. In-

jury, disability and quality of life after the 2009 earthquake in Padang, Indonesia: a prospective cohort study of adult survivors. *Glob Health Action*. 2012;**5**:1-11. doi: 10.3402/gha.v5i0.11816. [PubMed: 22629236].

 Hirshberg A, Holcomb JB, Mattox KL. Hospital trauma care in multiple-casualty incidents: a critical view. Ann Emerg Med. 2001;37(6):647-52. doi: 10.1067/mem.2001.115650. [PubMed: 11385336].