

# Surgical and Non-Surgical Management Strategies for Lateral Epicondylitis

Meisam Jafari Kafiabadi<sup>1</sup>, Amir Sabaghzadeh<sup>1</sup>, Farsad Biglari<sup>2</sup>, Amin Karami<sup>3</sup>, Mehrdad Sadighi<sup>1,4,\*</sup>, Adel Ebrahimpour<sup>4</sup>

<sup>1</sup> Assistant Professor, Department of Orthopedic Surgery, Shohada Tajrish Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>2</sup> Orthopedic Surgeon, Department of Orthopedic Surgery, Shohada Tajrish Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>3</sup> Resident, Department of Orthopedic Surgery, Shohada Tajrish Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>4</sup> Professor, Department of Orthopedic Surgery, Shohada Tajrish Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran

\* Corresponding author: Mehrdad Sadighi; Department of Orthopedic Surgery, Shohada Tajrish Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran.  
Tel: +98-9125754616, Email: mehrdad\_1330@yahoo.com

Received: 14 September 2020; Revised: 01 December 2020; Accepted: 12 January 2021

## Abstract

Lateral epicondylitis (LE) is one of the major causes of elbow pain. Despite being a self-limiting condition, its high incidence can cause a significant socioeconomic burden. Many treatment modalities have been proposed for the treatment, but the optimal strategy is still unknown. In this article, we discuss surgical and non-surgical strategies for the treatment of LE and address the research gaps.

**Keywords:** Exercise Therapy; Lateral Epicondylitis; Tennis Elbow; Treatment

**Citation:** Jafari Kafiabadi M, Sabaghzadeh A, Biglari F, Karami A, Sadighi M, Ebrahimpour A. **Surgical and Non-Surgical Management Strategies for Lateral Epicondylitis.** *J Orthop Spine Trauma* 2021; 7(1): 1-7.

## Background

Lateral epicondylitis (LE), otherwise known as “tennis elbow” is characterized by pain and tenderness over the lateral epicondyle of the humerus during the loading of the wrist extensor muscles. It affects 1-3 percent of the general population without any gender difference and usually presents between 35 and 54 years of age (1, 2). Individuals with repetitive motions of the forearm and/or fine motor skills, such as heavy manual laborers and tennis players, are at high risk of LE (3). Smoking and obesity are also associated with disease development (4).

LE tends to cause a significant economic and social burden due to the inability to perform routine activities and missed workdays. However, it is mostly a self-limited disease with symptom improvement in 70% to 90% of patients in weeks and complete resolution within a year (5, 6). There are numerous treatment options for this condition, but none are completely effective. In addition, in up to one-third of patients, the pain lasts more than a year which requires surgical treatment (7).

Here, we discuss surgical and non-surgical management strategies of LE based on available evidence and address research shortcomings in this field.

### Non-Surgical Management Strategies

#### I. Physiotherapy

Physiotherapy is almost always a part of any treatment regimen used for LE specially in the first 8 weeks (8, 9). Physiotherapy for LE consists of two main components: manual therapy and exercise therapy, which may or may not be used as a combination. When used in combination, there might be a better effect on clinical indicators such as pain-free grip (10). Manual therapy must be performed according to Mulligan's mobilization with movement (MWM). Ulnar-humeral lateral glide and radial head posteroanterior (PA) glide are two techniques used in the treatment of LE (10).

**I.A. Exercise Therapy:** The core principle of non-surgical management of LE is exercise. It has been proved to be beneficial as a part of a physical therapy regimen or separately (11). Some studies favor eccentric over concentric exercise (11-14). Eccentric exercise is performed by stretching of extensor compartment with an applied load. This type of exercise has been proved to be superior in comparison with other therapeutic modalities (12). Although useful, the optimum interval, duration, and intensity of this exercise have not been studied and there is a need for further investigations (14). In this section, we discuss common exercises.

**I.A1. Wrist Extensor Stretching:** This exercise consists of putting an elbow in full extension (with palms facing the ground) and wrist in 90° dorsiflexion while the opposite hand pushes the palm (Figure 1). The pressure should not induce pain and feeling a gentle stretch inside the forearm is enough. The stretched position should be kept for at least 15 seconds and the patient should perform the exercise for both sides 4 times a day (15).



**Figure 1.** Right wrist extensor muscles stretching




**I-A2. Wrist Flexor Stretching:** With the elbow in full extension (palms facing down) and wrist in 90° palmar flexion, the opposite hand is used to bend the hand toward the body until feeling a gentle stretch (Figure 2). Similar to the previous exercise, the position should be kept for at least 15 seconds and stretching should be repeated 4 times a day for both sides. This exercise could be done as a warm-up before activities involve gripping.




**Figure 2.** Right wrist flexor muscles stretching

**I-A3. Wrist Extensor Strengthening:** To perform an isotonic exercise of eccentric wrist extension (16), the patient should put the elbow in 90° flexion while supporting its weight by a stable surface with the wrist on the edge and palm facing the ground (Figure 3).



**Figure 3.** Right wrist eccentric extension exercise, starting position

Then the wrist is flexed upward to its maximum capacity, kept in position for 1 second, and then, slowly flexed downward (Figure 4).




**Figure 4.** Right wrist eccentric extension exercise, ending position

Depending on the severity of the pain felt by the patient during the exercise, it may be performed with or without a dumbbell (Figures 5 and 6).



**Figure 5.** Right wrist eccentric extension exercise with dumbbell, starting position

Each full wrist extension and flexion should take 5 to 10 seconds and the patient needs to repeat the exercise for both sides 10 times a day (11). In addition, the wrist extension can be performed with elbow in 180° extension using the same method to put more stress on extensors.



**Figure 6.** Right wrist eccentric extension exercise with dumbbell, ending position

**I-A4. Wrist Flexor Strengthening:** With the elbow in 90° flexion, its weight is supported with a stable surface by putting the wrist on the edge (palm facing up). Like the previous exercise, the wrist is flexed upward to its maximum capacity and kept in position for 1 second. Then, it is flexed downward slowly. To achieve better results, the patient can use a dumbbell if the pain is endurable. Each full wrist extension and flexion should take 5 to 10 seconds and the patient should repeat the exercise for both sides 10 times a day.

**I-A5. Twist Exercise:** In this exercise, the patient should grip a flexible bar by the affected hand, with the wrist in maximum extension, the elbow in 90° flexion, and the free end of the bar pointing up. The opposite hand grips the other end of the bar and twists the bar until the wrist reaches full flexion. Then, the elbows are fully extended in front of the body while both wrists are maintained in their previous positions. Finally, the affected wrist should be allowed to slowly become flexed by the force exerted by flexible bar untwisting (14).

## I-B. Physiotherapy Modalities

**I-B1. Shockwave Therapy:** Many studies have been conducted on shockwave therapy effectiveness in the past decade. In this therapeutic method, the shockwaves are aimed to the interstitial and extracellular responses that lead to tissue regeneration (17, 18). The absence of control groups and uniform treatment in these studies has yielded inconclusive results (19-21).

Recently, few studies with better design and similar treatment protocols have been published (22-24). In two meta-analyses comparing shockwave therapy with corticosteroid injection or ultrasonic waves, it has been proved that shockwave therapy is superior in terms of visual analog scale (VAS) and handgrip strength (25, 26). In a clinical trial, Ahadi et al. compared shockwave therapy with prolotherapy, ultrasound treatment, or placebo and found that shockwave therapy was significantly more effective than prolotherapy (27). In a recent study by Celik and Anafiroglu Kulunkoglu, photobiomodulation showed better efficacy in improving VAS and handgrip scores compared to shockwave therapy (28). Based on available evidence, shockwave therapy is not appropriate for acute LE, but is recommended when symptoms persist for more than 6 months or after the failure of other conservative treatments (8).

**I-B2. Taping:** Various taping techniques have been suggested for the treatment of LE (29). Although several studies have discussed the efficacy of taping procedures, a case-control design was not used and the evidence was low-quality (29, 30). Thus, it is unclear that the observed effects are due to the taping procedures or the course of the disease itself.

In a systematic review by George et al. in 2019, it was highlighted that with the current evidence, making a suggestion regarding efficiency of taping was not reasonable (30). In a meta-analysis, Zhong et al. investigated randomized controlled trials (RCTs) involving Kinesio taping (KT) (Figure 7) and concluded that this treatment improved VAS, handgrip, and the score of the Disabilities of the Arm, Shoulder and Hand (DASH) compared to placebo or physiotherapy (31). The main limitations of these RCTs were small sample size, various taping techniques, different treatment protocols, and short follow-up periods. Although the recent data have shown promising results, drawing a certain conclusion on this matter needs further studies with a uniform treatment strategy.



**Figure 7.** Kinesio taping (KT) to relieve tension of right extensor muscles

**I-B3. Acupuncture:** Rooted in traditional Chinese medicine, acupuncture has been used widely as a

treatment for pain relief in different musculoskeletal disorders (32-34). This method is also used in the treatment of LE. Although the definite mechanism of acupuncture is still unknown, it may increase local blood flow and cause neurotransmitter release which can facilitate healing and cause analgesia (35). In a recent meta-analysis of the RCTs by Morley et al., acupuncture has shown to be efficient and successful in relieving pain (36). Although many studies comparing acupuncture with different therapeutic methods have shown the benefits of acupuncture, most of them suffer from a small sample size (37-39).

There is a lack of data regarding acupuncture's long-term effects and possible side effects. Like many other procedures, acupuncture can be performed by different therapeutic instruments and on diverse locations. Due to the diversity of the therapeutic methods used by different studies, we cannot draw any conclusions regarding acupuncture's efficacy.

## II. Medications

### II-A. Nonsteroidal Anti-Inflammatory Drugs (NSAIDs):

Different oral NSAIDs have been used to treat LE. The use of tenoxicam, piroxicam, flurbiprofen, diclofenac, and naproxen has been effective in pain reduction but limited by gastrointestinal (GI) adverse effects (8). On the other hand, topical NSAIDs have favorable short-term effects and seem to possess no GI side effects (40). Although the NSAIDs may have short-term pain relief effects in LE, the long-term outcome is unclear and they cannot stop the disease progression (40, 41).

### II-B. Steroid Injection:

Introduced in the 1950s, corticosteroid injection is now one of the most widely-used treatments for LE (42, 43). In recent years, corticosteroid treatment has lost its popularity among practitioners due to the long-term inefficacy and side effects (44, 45). It has been associated with a rapid pain reduction in LE, but its effect on functional recovery is questionable. Although effective in the short term, corticosteroid injection is associated with relapses and poor long-term efficacy compared to other treatment modalities and even no treatment (46).

Corticosteroids exert their effects by inhibition of neuropeptides and cytokines. By doing so, they might also disrupt tendon healing by inhibiting migration and proliferation of cells and inducing the differentiation of the non-tenocytes (47-49). With such mechanisms, corticosteroid injection might also delay pain recovery in the long term by postponing tendon healing. In an RCT, Smidt et al. reported that corticosteroid injection even showed poor results compared to the "wait and see" approach (50). In another study, three or more corticosteroid injections were associated with the failure of surgical treatment (51). Recent researches have linked corticosteroid injection with tendon calcification and rupture, as well as decreased muscle thickness (52-54). It might also cause hypopigmentation and atrophy in the skin (55). Although it is a rapid solution to the patient's pain and poor function, we suggest that corticosteroid injection should be used with caution after considering other treatment options.

### II-C. Biological Therapeutics:

A new alternative treatment for LE is the injection of biological therapeutics. In this method, biological solutions such as autologous blood (AB) and platelet-rich plasma (PRP) are injected into the affected side to reduce pain.

For AB injection, 2 ml of AB is collected from the patient and injected proximal to the lateral epicondyle along the

supracondylar ridge in combination with lidocaine or bupivacaine (56). The needle should advance into the inferior surface of the extensor carpi radialis brevis muscle. These therapeutics contain growth factors that are beneficial for soft tissue healing. Moreover, an inflammatory reaction might be involved in the therapeutic effects observed with this treatment (57).

PRP is one of the biological products that have been intensively studied in the past few years. It is a derivate of AB which contains a more concentrated amount of platelet. There is no standard technique for the preparation of PRP. It can be prepared by centrifuging 15-20 ml of AB and could be injected in combination with different substances (56).

Consensus data suggest that AB and PRP are both superior to corticosteroid injection in terms of side effects and long-term effects, but corticosteroid injection has shown better outcomes in the short term (46, 58-60). A systematic review by Tang et al. suggests that PRP is superior to AB in terms of side effects (59). Various meta-analyses have been performed on RCTs addressing biological therapeutics injection (49, 58-61). Only one of them investigated PRP efficacy versus placebo treatment, and surprisingly, it found that there was no difference between them in terms of short- or long-term effects (61). This paradoxical finding might be due to the selection of saline as a placebo injection by many studies (62-64). As it was previously mentioned, LE is a self-limiting condition. Hence, it is of importance to compare each treatment with a placebo rather than other types of treatments.

To draw conclusions about biological therapeutics, we need studies that address different variables such as volume of injections, number of injections, injection intervals, and follow-up periods.

**II-D. Botulinum Injection:** Several studies have reported that botulinum injection can reduce pain in LE (65, 66). However, its application is associated with weakness. In a meta-analysis by Krogh et al., botulinum injection showed to be more effective than placebo injection, but less effective than corticosteroid injection in the short term and was associated with more grip weakness. This study also suggested that the duration of botulinum effect in LE was about 16 weeks (53). Based on available evidence, more studies with better study design and botulinum toxin dosage are required.

**II-E. Saline Injection:** Although considered as a placebo treatment, it has been recently postulated that saline injection might hold clinical and therapeutic effects in some musculoskeletal conditions such as LE. In a meta-analysis by Gao et al., the saline injection has shown therapeutic effects comparable to corticosteroid, PRP, autologous conditioned plasma (ACP), and botulinum injection (62). It has also successfully improved pain and functional scores (64). Whether these effects are a placebo effect or a property of the saline solution is unclear (62).

### Surgical Management Strategies

Surgery is the last resort for the treatment of LE. There are three main surgical approaches for this condition: open, arthroscopic, and percutaneous. Since the introduction of open surgery for the treatment of LE, this approach has evolved profoundly. The denervation of lateral epicondyle was one of the first procedures introduced by Wilhelm and Gieseler (67). Boyd and Mcleod introduced an open release procedure that includes removal of the annular ligament's proximal part and release of the epicondylar enthesis (68). Their technique was soon replaced by the Nirschl technique (69).

The main superiority of arthroscopic approaches in the treatment of LE is the ability to see intra-articular abnormalities, quicker recovery, and faster return to work (70). Although the percutaneous approach is not able to examine intra-articular abnormalities, it provides quicker recovery and better patient satisfaction. There are reports suggesting the same efficacy for the percutaneous approach compared to the open and arthroscopic methods (71). However, in one study, it was reported that open and arthroscopic approaches might result in higher DASH scores (72).

Radiofrequency microtenotomy is another method that has been successfully used in the treatment of LE. Results from comparative studies performed on radiofrequency microtenotomy and other surgical interventions have shown similar efficacy (73).

In recent years, promising results have been published on application of percutaneous ultrasonic tenotomy (72). This method has been used in a variety of orthopedic disorders and the majority of studies have shown satisfactory application of this procedure (74). Only one study compared this novel method with the previously-existing methods and reported a similar efficacy (75). However, it failed to report details of surgical procedures and had a very small sample size. Thus, the superiority of this method is yet to be determined.

Reports suggest a less than 10% failure rate for LE surgery (76). Various causes have been proposed for the failure of LE surgery. The most common cause of failure is patient non-compliance and inadequate rehabilitation (77). Another cause of surgical failure might be a concomitant orthopedic condition or inaccurate diagnosis (77). Radial tunnel syndrome (RTS) is one of the main causes of inaccurate diagnosis. In one study, up to 30% of patients diagnosed with LE were suffering from entrapment of the posterior interosseous nerve (78). Elbow instability and intra-articular complications such as synovial plica, synovial fistula, and osteoarthritis (OA) might also contribute to surgical failure and poor surgical outcomes. In 11% to 69% of patients affected by LE, intra-articular pathologies have been found (76). Surgical treatment in these cases usually results in worsening of symptoms or might contribute to the development of a different symptom complex (77).

### Conclusion

The treatment of LE is based on exercise therapy. There is no consensus regarding the next step in unresponsive patients due to the lack of evidence. PRP and shockwave therapy have shown promising results in comparison with other treatments. None of the existing surgical procedures have shown absolute superiority over each other. Although promising results have been observed with percutaneous ultrasonic tenotomy, further investigations are necessary.

### Conflict of Interest

The authors declare no conflict of interest in this study.

### Acknowledgments

None.

### References

1. Zwerus EL, Somford MP, Maissan F, Heisen J, Eygendaal D, van den Bekerom MP. Physical examination of the elbow, what is the evidence? A systematic literature review. *Br J*

- Sports Med.* 2018;52(19):1253-60. doi: [10.1136/bjsports-2016-096712](https://doi.org/10.1136/bjsports-2016-096712). [PubMed: [28249855](https://pubmed.ncbi.nlm.nih.gov/28249855/)].
2. Shiri R, Viikari-Juntura E, Varonen H, Heliovaara M. Prevalence and determinants of lateral and medial epicondylitis: A population study. *Am J Epidemiol.* 2006;164(11):1065-74. doi: [10.1093/aje/kwj325](https://doi.org/10.1093/aje/kwj325). [PubMed: [16968862](https://pubmed.ncbi.nlm.nih.gov/16968862/)].
  3. Haahr JP, Andersen JH. Physical and psychosocial risk factors for lateral epicondylitis: A population based case-referent study. *Occup Environ Med.* 2003;60(5):322-9. doi: [10.1136/oem.60.5.322](https://doi.org/10.1136/oem.60.5.322). [PubMed: [12709516](https://pubmed.ncbi.nlm.nih.gov/12709516/)]. [PubMed Central: [PMC1740535](https://pubmed.ncbi.nlm.nih.gov/PMC1740535/)].
  4. Vaquero-Picado A, Barco R, Antuna SA. Lateral epicondylitis of the elbow. *EFORT Open Rev.* 2016;1(11):391-7. doi: [10.1302/2058-5241.1.000049](https://doi.org/10.1302/2058-5241.1.000049). [PubMed: [28461918](https://pubmed.ncbi.nlm.nih.gov/28461918/)]. [PubMed Central: [PMC5367546](https://pubmed.ncbi.nlm.nih.gov/PMC5367546/)].
  5. Bisset L, Smidt N, Van der Windt DA, Bouter LM, Jull G, Brooks P, et al. Conservative treatments for tennis elbow do subgroups of patients respond differently? *Rheumatology (Oxford).* 2007;46(10):1601-5. doi: [10.1093/rheumatology/kem192](https://doi.org/10.1093/rheumatology/kem192). [PubMed: [17890274](https://pubmed.ncbi.nlm.nih.gov/17890274/)].
  6. Smidt N, Lewis M, Van der Windt DA, Hay EM, Bouter LM, Croft P. Lateral epicondylitis in general practice: Course and prognostic indicators of outcome. *J Rheumatol.* 2006;33(10):2053-9. [PubMed: [16881095](https://pubmed.ncbi.nlm.nih.gov/16881095/)].
  7. Meunier M. Lateral Epicondylitis/Extensor Tendon Injury. *Clin Sports Med.* 2020;39(3):657-60. doi: [10.1016/j.csm.2020.03.001](https://doi.org/10.1016/j.csm.2020.03.001). [PubMed: [32446581](https://pubmed.ncbi.nlm.nih.gov/32446581/)].
  8. Ma KL, Wang HQ. Management of lateral epicondylitis: A narrative literature review. *Pain Res Manag.* 2020;2020:6965381. doi: [10.1155/2020/6965381](https://doi.org/10.1155/2020/6965381). [PubMed: [32454922](https://pubmed.ncbi.nlm.nih.gov/32454922/)]. [PubMed Central: [PMC7222600](https://pubmed.ncbi.nlm.nih.gov/PMC7222600/)].
  9. Weber C, Thai V, Neuheuser K, Groover K, Christ O. Efficacy of physical therapy for the treatment of lateral epicondylitis: A meta-analysis. *BMC Musculoskelet Disord.* 2015;16:223. doi: [10.1186/s12891-015-0665-4](https://doi.org/10.1186/s12891-015-0665-4). [PubMed: [26303397](https://pubmed.ncbi.nlm.nih.gov/26303397/)]. [PubMed Central: [PMC4549077](https://pubmed.ncbi.nlm.nih.gov/PMC4549077/)].
  10. Bisset LM, Vicenzino B. Physiotherapy management of lateral epicondylalgia. *J Physiother.* 2015;61(4):174-81. doi: [10.1016/j.jphys.2015.07.015](https://doi.org/10.1016/j.jphys.2015.07.015). [PubMed: [26361816](https://pubmed.ncbi.nlm.nih.gov/26361816/)].
  11. Sevier TL, Stegink-Jansen CW. Astym treatment vs. eccentric exercise for lateral elbow tendinopathy: a randomized controlled clinical trial. *PeerJ.* 2015;3:e967. doi: [10.7717/peerj.967](https://doi.org/10.7717/peerj.967). [PubMed: [26038722](https://pubmed.ncbi.nlm.nih.gov/26038722/)]. [PubMed Central: [PMC4451036](https://pubmed.ncbi.nlm.nih.gov/PMC4451036/)].
  12. Cullinane FL, Boocock MG, Trevelyan FC. Is eccentric exercise an effective treatment for lateral epicondylitis? A systematic review. *Clin Rehabil.* 2014;28(1):3-19. doi: [10.1177/0269215513491974](https://doi.org/10.1177/0269215513491974). [PubMed: [23881334](https://pubmed.ncbi.nlm.nih.gov/23881334/)].
  13. Peterson M, Butler S, Eriksson M, Svardsudd K. A randomized controlled trial of eccentric vs. concentric graded exercise in chronic tennis elbow (lateral elbow tendinopathy). *Clin Rehabil.* 2014;28(9):862-72. doi: [10.1177/0269215514527595](https://doi.org/10.1177/0269215514527595). [PubMed: [24634444](https://pubmed.ncbi.nlm.nih.gov/24634444/)].
  14. Raman J, MacDermid JC, Grewal R. Effectiveness of different methods of resistance exercises in lateral epicondylitis—a systematic review. *J Hand Ther.* 2012;25(1):5-25. doi: [10.1016/j.jht.2011.09.001](https://doi.org/10.1016/j.jht.2011.09.001). [PubMed: [22075055](https://pubmed.ncbi.nlm.nih.gov/22075055/)].
  15. Martinez-Silvestrini JA, Newcomer KL, Gay RE, Schaefer MP, Kortebein P, Arendt KW. Chronic lateral epicondylitis: comparative effectiveness of a home exercise program including stretching alone versus stretching supplemented with eccentric or concentric strengthening. *J Hand Ther.* 2005;18(4):411-9, quiz. doi: [10.1197/j.jht.2005.07.007](https://doi.org/10.1197/j.jht.2005.07.007). [PubMed: [16271688](https://pubmed.ncbi.nlm.nih.gov/16271688/)].
  16. Tyler TF, Thomas GC, Nicholas SJ, McHugh MP. Addition of isolated wrist extensor eccentric exercise to standard treatment for chronic lateral epicondylitis: a prospective randomized trial. *J Shoulder Elbow Surg.* 2010;19(6):917-22. doi: [10.1016/j.jse.2010.04.041](https://doi.org/10.1016/j.jse.2010.04.041). [PubMed: [20579907](https://pubmed.ncbi.nlm.nih.gov/20579907/)].
  17. Ogden JA, Toth-Kischkat A, Schultheiss R. Principles of shock wave therapy. *Clin Orthop Relat Res.* 2001;(387):8-17. doi: [10.1097/00003086-200106000-00003](https://doi.org/10.1097/00003086-200106000-00003). [PubMed: [11400898](https://pubmed.ncbi.nlm.nih.gov/11400898/)].
  18. Siebert WE, Buch CM. Extracorporeal shock waves in orthopaedics. Berlin, Germany: Springer Berlin Heidelberg; 2012.
  19. Crevenna R, Mickel M, Keilani M. Extracorporeal shock wave therapy in the supportive care and rehabilitation of cancer patients. *Support Care Cancer.* 2019;27(11):4039-41. doi: [10.1007/s00520-019-05046-y](https://doi.org/10.1007/s00520-019-05046-y). [PubMed: [31446484](https://pubmed.ncbi.nlm.nih.gov/31446484/)]. [PubMed Central: [PMC6803573](https://pubmed.ncbi.nlm.nih.gov/PMC6803573/)].
  20. Razavipour M, Azar MS, Kariminasab MH, Gaffari S, Fazli M. The short term effects of shock-wave therapy for tennis elbow: A clinical trial study. *Acta Inform Med.* 2018;26(1):54-6. doi: [10.5455/aim.2018.26.54-56](https://doi.org/10.5455/aim.2018.26.54-56). [PubMed: [29719315](https://pubmed.ncbi.nlm.nih.gov/29719315/)]. [PubMed Central: [PMC5869227](https://pubmed.ncbi.nlm.nih.gov/PMC5869227/)].
  21. Testa G, Vescio A, Perez S, Petrantonio V, Mazzarella G, Costarella L, et al. Functional outcome at short and middle term of the extracorporeal shockwave therapy treatment in lateral epicondylitis: A case-series study. *J Clin Med.* 2020;9(3):633. doi: [10.3390/jcm9030633](https://doi.org/10.3390/jcm9030633). [PubMed: [32120893](https://pubmed.ncbi.nlm.nih.gov/32120893/)]. [PubMed Central: [PMC7141112](https://pubmed.ncbi.nlm.nih.gov/PMC7141112/)].
  22. Dedes V, Tzirogiannis K, Polikandrioti M, Dede AM, Mitseas A, Panoutsopoulos GI. Comparison of radial extracorporeal shockwave therapy with ultrasound therapy in patients with lateral epicondylitis. *J Med Ultrason (2001).* 2020;47(2):319-25. doi: [10.1007/s10396-019-01002-9](https://doi.org/10.1007/s10396-019-01002-9). [PubMed: [31912320](https://pubmed.ncbi.nlm.nih.gov/31912320/)].
  23. Guler NS, Sargin S, Sahin N. Efficacy of extracorporeal shockwave therapy in patients with lateral epicondylitis: A randomized, placebo-controlled, double-blind clinical trial. *North Clin Istanb.* 2018;5(4):314-8. doi: [10.14744/nci.2017.82435](https://doi.org/10.14744/nci.2017.82435). [PubMed: [30859161](https://pubmed.ncbi.nlm.nih.gov/30859161/)]. [PubMed Central: [PMC6371991](https://pubmed.ncbi.nlm.nih.gov/PMC6371991/)].
  24. Mastej S, Pop T, Bejer A, Plocki J, Kotela I. Comparison of the effectiveness of shockwave therapy with selected physical therapy procedures in patients with tennis elbow syndrome. *Ortop Traumatol Rehabil.* 2018;20(4):301-11. doi: [10.5604/01.3001.0012.6464](https://doi.org/10.5604/01.3001.0012.6464). [PubMed: [30648651](https://pubmed.ncbi.nlm.nih.gov/30648651/)].
  25. Xiong Y, Xue H, Zhou W, Sun Y, Liu Y, Wu Q, et al. Shock-wave therapy versus corticosteroid injection on lateral epicondylitis: A meta-analysis of randomized controlled trials. *Phys Sportsmed.* 2019;47(3):284-9. doi: [10.1080/00913847.2019.1599587](https://doi.org/10.1080/00913847.2019.1599587). [PubMed: [30951399](https://pubmed.ncbi.nlm.nih.gov/30951399/)].
  26. Yan C, Xiong Y, Chen L, Endo Y, Hu L, Liu M, et al. A comparative study of the efficacy of ultrasonics and extracorporeal shock wave in the treatment of tennis elbow: A meta-analysis of randomized controlled trials. *J Orthop Surg Res.* 2019;14(1):248. doi: [10.1186/s13018-019-1290-y](https://doi.org/10.1186/s13018-019-1290-y). [PubMed: [31387611](https://pubmed.ncbi.nlm.nih.gov/31387611/)]. [PubMed Central: [PMC6683364](https://pubmed.ncbi.nlm.nih.gov/PMC6683364/)].
  27. Ahadi T, Esmaili Jamkarani M., Raissi GR, Mansoori K, Emami Razavi SZ, Sajadi S. Prolotherapy vs radial extracorporeal shock wave therapy in the short-term treatment of lateral epicondylitis: A randomized clinical trial. *Pain Med.* 2019;20(9):1745-9. doi: [10.1093/pm/pty303](https://doi.org/10.1093/pm/pty303). [PubMed: [30698771](https://pubmed.ncbi.nlm.nih.gov/30698771/)].
  28. Celik D, Anaforglu Kulunkoglu B. Photobiomodulation Therapy Versus extracorporeal shock wave therapy in the treatment of lateral epicondylitis. *Photobiomodul Photomed Laser Surg.* 2019;37(5):269-75. doi: [10.1089/photob.2018.4533](https://doi.org/10.1089/photob.2018.4533). [PubMed: [31084562](https://pubmed.ncbi.nlm.nih.gov/31084562/)].
  29. Dilek B, Batmaz I, Sariyildiz MA, Sahin E, Ilter L, Gulbahar S, et al. Kinesio taping in patients with lateral epicondylitis. *J Back Musculoskelet Rehabil.* 2016;29(4):853-8. doi: [10.3233/BMR-160701](https://doi.org/10.3233/BMR-160701). [PubMed: [27232082](https://pubmed.ncbi.nlm.nih.gov/27232082/)].
  30. George CE, Heales LJ, Stanton R, Wintour SA, Kean CO. Sticking to the facts: A systematic review of the effects of therapeutic

- tape in lateral epicondylalgia. *Phys Ther Sport*. 2019;40:117-27. doi: [10.1016/j.pts.2019.08.011](https://doi.org/10.1016/j.pts.2019.08.011). [PubMed: [31518778](https://pubmed.ncbi.nlm.nih.gov/31518778/)].
31. Zhong Y, Zheng C, Zheng J, Xu S. Kinesio tape reduces pain in patients with lateral epicondylitis: A meta-analysis of randomized controlled trials. *Int J Surg*. 2020;76:190-9. doi: [10.1016/j.ijsu.2020.02.044](https://doi.org/10.1016/j.ijsu.2020.02.044). [PubMed: [32165280](https://pubmed.ncbi.nlm.nih.gov/32165280/)].
  32. Kaptchuk TJ. Acupuncture: theory, efficacy, and practice. *Ann Intern Med*. 2002;136(5):374-83. doi: [10.7326/0003-4819-136-5-200203050-00010](https://doi.org/10.7326/0003-4819-136-5-200203050-00010). [PubMed: [11874310](https://pubmed.ncbi.nlm.nih.gov/11874310/)].
  33. Sierpina VS, Frenkel MA. Acupuncture: A clinical review. *South Med J*. 2005;98(3):330-7. doi: [10.1097/01.SMJ.0000140834.30654.0F](https://doi.org/10.1097/01.SMJ.0000140834.30654.0F). [PubMed: [15813160](https://pubmed.ncbi.nlm.nih.gov/15813160/)].
  34. Eisenberg DM, Kessler RC, Foster C, Norlock FE, Calkins DR, Delbanco TL. Unconventional medicine in the United States. Prevalence, costs, and patterns of use. *N Engl J Med*. 1993;328(4):246-52. doi: [10.1056/NEJM199301283280406](https://doi.org/10.1056/NEJM199301283280406). [PubMed: [8418405](https://pubmed.ncbi.nlm.nih.gov/8418405/)].
  35. Cheng KJ. Neurobiological mechanisms of acupuncture for some common illnesses: A clinician's perspective. *J Acupunct Meridian Stud*. 2014;7(3):105-14. doi: [10.1016/j.jams.2013.07.008](https://doi.org/10.1016/j.jams.2013.07.008). [PubMed: [24929454](https://pubmed.ncbi.nlm.nih.gov/24929454/)].
  36. Morley S, Eccleston C, Williams A. Systematic review and meta-analysis of randomized controlled trials of cognitive behaviour therapy and behaviour therapy for chronic pain in adults, excluding headache. *Pain*. 1999;80(1-2):1-13. doi: [10.1016/s0304-3959\(98\)00255-3](https://doi.org/10.1016/s0304-3959(98)00255-3). [PubMed: [10204712](https://pubmed.ncbi.nlm.nih.gov/10204712/)].
  37. Haker E, Lundeberg T. Acupuncture treatment in epicondylalgia: A comparative study of two acupuncture techniques. *Clin J Pain*. 1990;6(3):221-6. doi: [10.1097/00002508-199009000-00009](https://doi.org/10.1097/00002508-199009000-00009). [PubMed: [2135016](https://pubmed.ncbi.nlm.nih.gov/2135016/)].
  38. Irnich D, Karg H, Behrens N, Lang M, Schreiber A, Kroling P. Controlled trial on point specificity of acupuncture in the treatment of lateral Epicondylitis (Tennis Elbow). *Physikalische Medizin, Rehabilitationsmedizin, Kurortmedizin*. 2003;13(4):215-9. doi: [10.1055/s-2003-41368](https://doi.org/10.1055/s-2003-41368).
  39. Fink M, Wolkenstein E, Karst M, Gehrke A. Acupuncture in chronic epicondylitis: A randomized controlled trial. *Rheumatology (Oxford)*. 2002;41(2):205-9. doi: [10.1093/rheumatology/41.2.205](https://doi.org/10.1093/rheumatology/41.2.205). [PubMed: [11886971](https://pubmed.ncbi.nlm.nih.gov/11886971/)].
  40. Pattanittum P, Turner T, Green S, Buchbinder R. Non-steroidal anti-inflammatory drugs (NSAIDs) for treating lateral elbow pain in adults. *Cochrane Database Syst Rev*. 2013;(5):CD003686. doi: [10.1002/14651858.CD003686.pub2](https://doi.org/10.1002/14651858.CD003686.pub2). [PubMed: [23728646](https://pubmed.ncbi.nlm.nih.gov/23728646/)]. [PubMed Central: [PMC7173751](https://pubmed.ncbi.nlm.nih.gov/PMC7173751/)].
  41. Abat F, Alfredson H, Cucchiari M, Madry H, Marmotti A, Mouton C, et al. Current trends in tendinopathy: consensus of the ESSKA basic science committee. Part II: Treatment options. *J Exp Orthop*. 2018;5(1):38. doi: [10.1186/s40634-018-0145-5](https://doi.org/10.1186/s40634-018-0145-5). [PubMed: [30251203](https://pubmed.ncbi.nlm.nih.gov/30251203/)]. [PubMed Central: [PMC6153202](https://pubmed.ncbi.nlm.nih.gov/PMC6153202/)].
  42. Titchener AG, Booker SJ, Bhamber NS, Tambe AA, Clark DI. Corticosteroid and platelet-rich plasma injection therapy in tennis elbow (lateral epicondylalgia): A survey of current U.K. specialist practice and a call for clinical guidelines. *Br J Sports Med*. 2015;49(21):1410-3. doi: [10.1136/bjsports-2013-092674](https://doi.org/10.1136/bjsports-2013-092674). [PubMed: [24195919](https://pubmed.ncbi.nlm.nih.gov/24195919/)].
  43. Cyriax J, Troisier O. Hydrocortone and soft-tissue lesions. *Br Med J*. 1953;2(4843):966-8. doi: [10.1136/bmj.2.4843.966](https://doi.org/10.1136/bmj.2.4843.966). [PubMed: [13094084](https://pubmed.ncbi.nlm.nih.gov/13094084/)]. [PubMed Central: [PMC2029934](https://pubmed.ncbi.nlm.nih.gov/PMC2029934/)].
  44. Coombes BK, Bisset L, Brooks P, Khan A, Vicenzino B. Effect of corticosteroid injection, physiotherapy, or both on clinical outcomes in patients with unilateral lateral epicondylalgia: A randomized controlled trial. *JAMA*. 2013;309(5):461-9. doi: [10.1001/jama.2013.129](https://doi.org/10.1001/jama.2013.129). [PubMed: [23385272](https://pubmed.ncbi.nlm.nih.gov/23385272/)].
  45. Coombes BK, Bisset L, Vicenzino B. Efficacy and safety of corticosteroid injections and other injections for management of tendinopathy: A systematic review of randomised controlled trials. *Lancet*. 2010;376(9754):1751-67. doi: [10.1016/S0140-6736\(10\)61160-9](https://doi.org/10.1016/S0140-6736(10)61160-9). [PubMed: [20970844](https://pubmed.ncbi.nlm.nih.gov/20970844/)].
  46. Houck DA, Kraeutler MJ, Thornton LB, McCarty EC, Bravman JT. Treatment of lateral epicondylitis with autologous blood, platelet-rich plasma, or corticosteroid injections: A systematic review of overlapping meta-analyses. *Orthop J Sports Med*. 2019;7(3):2325967119831052. doi: [10.1177/2325967119831052](https://doi.org/10.1177/2325967119831052). [PubMed: [30899764](https://pubmed.ncbi.nlm.nih.gov/30899764/)]. [PubMed Central: [PMC6419259](https://pubmed.ncbi.nlm.nih.gov/PMC6419259/)].
  47. Chou LC, Liou TH, Kuan YC, Huang YH, Chen HC. Autologous blood injection for treatment of lateral epicondylitis: A meta-analysis of randomized controlled trials. *Phys Ther Sport*. 2016;18:68-73. doi: [10.1016/j.pts.2015.06.002](https://doi.org/10.1016/j.pts.2015.06.002). [PubMed: [26616689](https://pubmed.ncbi.nlm.nih.gov/26616689/)].
  48. Qian X, Lin Q, Wei K, Hu B, Jing P, Wang J. Efficacy and safety of autologous blood products compared with corticosteroid injections in the treatment of lateral epicondylitis: A meta-analysis of randomized controlled trials. *PM R*. 2016;8(8):780-91. doi: [10.1016/j.pmrj.2016.02.008](https://doi.org/10.1016/j.pmrj.2016.02.008). [PubMed: [26968611](https://pubmed.ncbi.nlm.nih.gov/26968611/)].
  49. Mi B, Liu G, Zhou W, Lv H, Liu Y, Wu Q, et al. Platelet rich plasma versus steroid on lateral epicondylitis: Meta-analysis of randomized clinical trials. *Phys Sportsmed*. 2017;45(2):97-104. doi: [10.1080/00913847.2017.1297670](https://doi.org/10.1080/00913847.2017.1297670). [PubMed: [28276986](https://pubmed.ncbi.nlm.nih.gov/28276986/)].
  50. Smidt N, Van der Windt DA, Assendelft WJ, Deville WL, Korthals-de Bos IB, Bouter LM. Corticosteroid injections, physiotherapy, or a wait-and-see policy for lateral epicondylitis: A randomised controlled trial. *Lancet*. 2002;359(9307):657-62. doi: [10.1016/S0140-6736\(02\)07811-X](https://doi.org/10.1016/S0140-6736(02)07811-X). [PubMed: [11879861](https://pubmed.ncbi.nlm.nih.gov/11879861/)].
  51. Degen RM, Cancienne JM, Camp CL, Altchek DW, Dines JS, Werner BC. Three or more preoperative injections is the most significant risk factor for revision surgery after operative treatment of lateral epicondylitis: An analysis of 3863 patients. *J Shoulder Elbow Surg*. 2017;26(4):704-9. doi: [10.1016/j.jse.2016.10.022](https://doi.org/10.1016/j.jse.2016.10.022). [PubMed: [28094190](https://pubmed.ncbi.nlm.nih.gov/28094190/)].
  52. Park HB, Kam M, Gwark JY. Association of steroid injection with soft-tissue calcification in lateral epicondylitis. *J Shoulder Elbow Surg*. 2019;28(2):304-9. doi: [10.1016/j.jse.2018.10.009](https://doi.org/10.1016/j.jse.2018.10.009). [PubMed: [30658775](https://pubmed.ncbi.nlm.nih.gov/30658775/)].
  53. Krogh TP, Fredberg U, Stengaard-Pedersen K, Christensen R, Jensen P, Ellingsen T. Treatment of lateral epicondylitis with platelet-rich plasma, glucocorticoid, or saline: a randomized, double-blind, placebo-controlled trial. *Am J Sports Med*. 2013;41(3):625-35. doi: [10.1177/0363546512472975](https://doi.org/10.1177/0363546512472975). [PubMed: [23328738](https://pubmed.ncbi.nlm.nih.gov/23328738/)].
  54. Smith AG, Kosygan K, Williams H, Newman RJ. Common extensor tendon rupture following corticosteroid injection for lateral tendinosis of the elbow. *Br J Sports Med*. 1999;33(6):423-4. doi: [10.1136/bjsm.33.6.423](https://doi.org/10.1136/bjsm.33.6.423). [PubMed: [10597854](https://pubmed.ncbi.nlm.nih.gov/10597854/)]. [PubMed Central: [PMC1756217](https://pubmed.ncbi.nlm.nih.gov/PMC1756217/)].
  55. Weinhammer AP, Shields BE, Keenan T. Intralesional corticosteroid-induced hypopigmentation and atrophy. *Dermatol Online J*. 2020;26(1):13030. [PubMed: [32155037](https://pubmed.ncbi.nlm.nih.gov/32155037/)].
  56. Calandruccio JH, Steiner MM. Autologous blood and platelet-rich plasma injections for treatment of lateral epicondylitis. *Orthop Clin North Am*. 2017;48(3):351-7. doi: [10.1016/j.ocl.2017.03.011](https://doi.org/10.1016/j.ocl.2017.03.011). [PubMed: [28577784](https://pubmed.ncbi.nlm.nih.gov/28577784/)].
  57. Tarpada SP, Morris MT, Lian J, Rashidi S. Current advances in the treatment of medial and lateral epicondylitis. *J Orthop*. 2018;15(1):107-10. doi: [10.1016/j.jor.2018.01.040](https://doi.org/10.1016/j.jor.2018.01.040). [PubMed: [29657450](https://pubmed.ncbi.nlm.nih.gov/29657450/)]. [PubMed Central: [PMC5895908](https://pubmed.ncbi.nlm.nih.gov/PMC5895908/)].
  58. Xu Q, Chen J, Cheng L. Comparison of platelet rich plasma and corticosteroids in the management of lateral epicondylitis: A meta-analysis of randomized controlled trials. *Int J Surg*. 2019;67:37-46. doi: [10.1016/j.ijsu.2019.05.003](https://doi.org/10.1016/j.ijsu.2019.05.003). [PubMed: [31128316](https://pubmed.ncbi.nlm.nih.gov/31128316/)].
  59. Tang S, Wang X, Wu P, Wu P, Yang J, Du Z, et al. Platelet-rich plasma Vs autologous blood Vs corticosteroid injections in the treatment of lateral epicondylitis: A systematic review,

- pairwise and network meta-analysis of randomized controlled trials. *PM R*. 2020;12(4):397-409. doi: [10.1002/pmrj.12287](https://doi.org/10.1002/pmrj.12287). [PubMed: [31736257](https://pubmed.ncbi.nlm.nih.gov/31736257/)]. [PubMed Central: [PMC7187193](https://pubmed.ncbi.nlm.nih.gov/PMC7187193/)].
60. Li A, Wang H, Yu Z, Zhang G, Feng S, Liu L, et al. Platelet-rich plasma vs corticosteroids for elbow epicondylitis: A systematic review and meta-analysis. *Medicine (Baltimore)*. 2019;98(51):e18358. doi: [10.1097/MD.00000000000018358](https://doi.org/10.1097/MD.00000000000018358). [PubMed: [31860992](https://pubmed.ncbi.nlm.nih.gov/31860992/)]. [PubMed Central: [PMC6940118](https://pubmed.ncbi.nlm.nih.gov/PMC6940118/)].
  61. Simental-Mendia M, Vilchez-Cavazos F, Alvarez-Villalobos N, Blazquez-Saldana J, Pena-Martinez V, Villarreal-Villarreal G, et al. Clinical efficacy of platelet-rich plasma in the treatment of lateral epicondylitis: A systematic review and meta-analysis of randomized placebo-controlled clinical trials. *Clin Rheumatol*. 2020;39(8):2255-65. doi: [10.1007/s10067-020-05000-y](https://doi.org/10.1007/s10067-020-05000-y). [PubMed: [32103373](https://pubmed.ncbi.nlm.nih.gov/32103373/)].
  62. Gao B, Dwivedi S, DeFroda S, Bokshan S, Ready LV, Cole BJ, et al. The therapeutic benefits of saline solution injection for lateral epicondylitis: A meta-analysis of randomized controlled trials comparing saline injections with nonsurgical injection therapies. *Arthroscopy*. 2019;35(6):1847-59. doi: [10.1016/j.arthro.2019.02.051](https://doi.org/10.1016/j.arthro.2019.02.051). [PubMed: [31072722](https://pubmed.ncbi.nlm.nih.gov/31072722/)].
  63. Saltzman BM, Leroux T, Meyer MA, Basques BA, Chahal J, Bach BR Jr, et al. The therapeutic effect of intra-articular normal saline injections for knee osteoarthritis: A meta-analysis of evidence level 1 studies. *Am J Sports Med*. 2017;45(11):2647-53. doi: [10.1177/0363546516680607](https://doi.org/10.1177/0363546516680607). [PubMed: [28027657](https://pubmed.ncbi.nlm.nih.gov/28027657/)].
  64. Acosta-Olivo CA, Millan-Alanis JM, Simental-Mendia LE, Alvarez-Villalobos N, Vilchez-Cavazos F, Pena-Martinez VM, et al. Effect of normal saline injections on lateral epicondylitis symptoms: A systematic review and meta-analysis of randomized clinical trials. *Am J Sports Med*. 2020;48(12):3094-102. doi: [10.1177/0363546519899644](https://doi.org/10.1177/0363546519899644). [PubMed: [32045280](https://pubmed.ncbi.nlm.nih.gov/32045280/)].
  65. Galvin R, Callaghan C, Chan WS, Dimitrov BD, Fahey T. Injection of botulinum toxin for treatment of chronic lateral epicondylitis: Systematic review and meta-analysis. *Semin Arthritis Rheum*. 2011;40(6):585-7. doi: [10.1016/j.semarthrit.2011.01.003](https://doi.org/10.1016/j.semarthrit.2011.01.003). [PubMed: [21397930](https://pubmed.ncbi.nlm.nih.gov/21397930/)].
  66. Jabbari B, Machado D. Treatment of refractory pain with botulinum toxins-an evidence-based review. *Pain Med*. 2011;12(11):1594-606. doi: [10.1111/j.1526-4637.2011.01245.x](https://doi.org/10.1111/j.1526-4637.2011.01245.x). [PubMed: [21958302](https://pubmed.ncbi.nlm.nih.gov/21958302/)].
  67. Wilhelm A, Gieseler H. Treatment of radiohumeral epicondylitis by denervation. *Chirurg*. 1962;33:118-22. [PubMed: [14007004](https://pubmed.ncbi.nlm.nih.gov/14007004/)].
  68. Boyd HB, McLeod AC Jr. Tennis elbow. *J Bone Joint Surg Am*. 1973;55(6):1183-7. [PubMed: [4585946](https://pubmed.ncbi.nlm.nih.gov/4585946/)].
  69. Nirschl RP, Pettrone FA. Tennis elbow. The surgical treatment of lateral epicondylitis. *J Bone Joint Surg Am*. 1979;61(6A):832-9. [PubMed: [479229](https://pubmed.ncbi.nlm.nih.gov/479229/)].
  70. Baker CL Jr, Murphy KP, Gottlob CA, Curd DT. Arthroscopic classification and treatment of lateral epicondylitis: Two-year clinical results. *J Shoulder Elbow Surg*. 2000;9(6):475-82. doi: [10.1067/mse.2000.108533](https://doi.org/10.1067/mse.2000.108533). [PubMed: [11155299](https://pubmed.ncbi.nlm.nih.gov/11155299/)].
  71. Burn MB, Mitchell RJ, Liberman SR, Lintner DM, Harris JD, McCulloch PC. Open, arthroscopic, and percutaneous surgical treatment of lateral epicondylitis: A systematic review. *Hand (NY)*. 2018;13(3):264-74. doi: [10.1177/1558944717701244](https://doi.org/10.1177/1558944717701244). [PubMed: [28720043](https://pubmed.ncbi.nlm.nih.gov/28720043/)]. [PubMed Central: [PMC5987981](https://pubmed.ncbi.nlm.nih.gov/PMC5987981/)].
  72. Bateman M, Littlewood C, Rawson B, Tambe AA. Surgery for tennis elbow: A systematic review. *Shoulder Elbow*. 2019;11(1):35-44. doi: [10.1177/1758573217745041](https://doi.org/10.1177/1758573217745041). [PubMed: [30719096](https://pubmed.ncbi.nlm.nih.gov/30719096/)]. [PubMed: [PMC6348580](https://pubmed.ncbi.nlm.nih.gov/PMC6348580/)].
  73. Lee JH, Park I, Hyun HS, Shin SJ. A comparison of radiofrequency-based microtenotomy and arthroscopic release of the extensor carpi radialis brevis tendon in recalcitrant lateral epicondylitis: A prospective randomized controlled study. *Arthroscopy*. 2018;34(5):1439-46. doi: [10.1016/j.arthro.2017.11.029](https://doi.org/10.1016/j.arthro.2017.11.029). [PubMed: [29366739](https://pubmed.ncbi.nlm.nih.gov/29366739/)].
  74. Jacobson JA, Kim SM, Brigido MK. Ultrasound-guided percutaneous tenotomy. *Semin Musculoskelet Radiol*. 2016;20(5):414-21. doi: [10.1055/s-0036-1597545](https://doi.org/10.1055/s-0036-1597545). [PubMed: [28002863](https://pubmed.ncbi.nlm.nih.gov/28002863/)].
  75. Althawi F, Li X, Demarest B, Forney MC. Percutaneous ultrasonic tenotomy with the TX-1 device versus surgical tenotomy for the treatment of common extensor tendinosis. *Skeletal Radiol*. 2021;50(1):115-24. doi: [10.1007/s00256-020-03540-7](https://doi.org/10.1007/s00256-020-03540-7). [PubMed: [32647907](https://pubmed.ncbi.nlm.nih.gov/32647907/)].
  76. Pomerantz ML. Complications of lateral epicondylar release. *Orthop Clin North Am*. 2016;47(2):445-69. doi: [10.1016/j.ocl.2015.10.002](https://doi.org/10.1016/j.ocl.2015.10.002). [PubMed: [26772953](https://pubmed.ncbi.nlm.nih.gov/26772953/)].
  77. Morrey BF. Reoperation for failed surgical treatment of refractory lateral epicondylitis. *J Shoulder Elbow Surg*. 1992;1(1):47-55. doi: [10.1016/S1058-2746\(09\)80016-5](https://doi.org/10.1016/S1058-2746(09)80016-5). [PubMed: [22958970](https://pubmed.ncbi.nlm.nih.gov/22958970/)].
  78. Jaloavaara P, Lindholm RV. Decompression of the posterior interosseous nerve for tennis elbow. *Arch Orthop Trauma Surg*. 1989;108(4):243-5. doi: [10.1007/BF00936209](https://doi.org/10.1007/BF00936209). [PubMed: [2774878](https://pubmed.ncbi.nlm.nih.gov/2774878/)].